

# Partial-equilibrium concepts to model trace element uptake

DMITRII A. KULIK<sup>1\*</sup>, BRUNO THIEN<sup>1</sup>, ENZO CURTI<sup>1</sup>

<sup>1</sup>Paul Scherrer Institut, Villigen PSI, Switzerland,  
dmitrii.kulik@psi.ch (\* presenting author)

Uptake of trace elements into host minerals involves various mechanisms - from ion exchange and inner-sphere adsorption via surface entrapment or in-diffusion to structural incorporation upon precipitation or recrystallization. Even at long reaction time, their interplay may result in a metastable distribution of minor elements that deviates from complete aqueous- solid solution thermodynamic equilibrium. Uptake processes may proceed at widely different rates, e.g. fast (minutes to hours) for ion exchange and slow for recrystallization (days to years), with important consequences for the reactive transport of contaminants. In this work, we explored how the rate-dependent uptake and metastability can be implemented in geochemical models using the partial equilibrium principle.

Thermodynamic models cannot be directly related to particular molecular mechanisms or their locations. Yet, some information can be retained in the model by introducing (i) multiple metastability constraints; (ii) mineral surface areas and site densities; and (iii) a set of 'layered', intrinsically metastable solid phases. In particular, the additional time-step-dependent metastability constraints [1] can be imposed on the amounts of solid solution end members. Because the specific free energy of the solid-aqueous interface is usually positive, solids with large specific surface area are metastable relative to coarse-crystalline counterparts. The surface free energy decreases due to adsorption, the extent of which is limited by the site density and the solid surface area. The limited adsorption can be modeled using the activity coefficient terms [2] or, more commonly, by adding the mass balances for surface- or permanent charge sites [3,4].

In usual aqueous - solid solution (AqSS) models, metastability constraints can only be set on bulk amounts of host and trace end members. In the new aqueous - layered solid (AqLS) model, 'seed' grains with defined specific surface, representing metastable cores of the solid, are enveloped by layers of the overgrown (or recrystallized) solid solution phase. This 'overgrowth' part has its own metastability constraints linked to the surface area of 'seed' phase. The 'adsorption solution' phase can be linked to the surface area of either the 'layer' (in AqALS) or the 'seed' phase (AqAS).

These partial equilibrium concepts, combined with different models of uptake kinetics [5], have been implemented in the GEM-Selektor code (<http://gems.web.psi.ch>) and tested in simulations of uptake into carbonates, sulfates, and goethite. Major trends in the experimental data were reproduced well in all cases. A more detailed time-dependent distribution of trace elements in AqLS and AqALS models could be reconciled with the inferred uptake mechanisms.

The research leading to these results has received funding from the European Atomic Energy Community's Seventh Framework Programme (FP7/2007-2011) under grant agreement n° 269688.

[1] Karpov et al. (2001) *Geochem. Internat.* **39**, 1108-1119. [2] Kulik (2006) *Interf. Sci. Technol.* **11**, 171-250. [3] Bradbury & Baeyens (2006) *Interf. Sci. Technol.* **11**, 518-538. [4] Kulik (2009) *RiMG* **70**, 125-180. [5] Thien et al. (2012), this conference.